#### International Journal of Mechanical Engineering and Technology (IJMET)

Volume 9, Issue 8, August 2018, pp. 1022–1028, Article ID: IJMET\_09\_08\_111 Available online at http://www.iaeme.com/ijmet/issues.asp?JType=IJMET&VType=9&IType=8 ISSN Print: 0976-6340 and ISSN Online: 0976-6359

© IAEME Publication



Scopus Indexed

# ANALYSIS OF VARIETY AND PLANTING TIME OF INDONESIAN SOYBEAN TOWARDS THE PRODUCTION RESULT TO MEET THE DEMAND

Nelly Budiharti

Industrial Engineering, National Institute of Technology (ITN), Malang

#### ING Wardana

Mechanical Engineering, Brawijaya University, Malang

#### ABSTRACT

To meet the high demand of soybean from domestic soybean production it is necessary to increase the planting time. For years, soybean farmers only plant once in a year that is in June or July. This research was conducted by experiment using 5 superior varieties planted in March, April and May. Data processing and analysis use experimental design of factorial design with random model of 95% confidence level. The results obtained that soybean varieties and planting time does not affect the production results, whether the interaction between soybean varieties and planting time greatly affects the production.

**Keywords**: Indonesian Soybean Varieties, Planting Time, Interaction, Production Results

**Cite this Article:** Nelly Budiharti and ING Wardana, Analysis of Variety and Planting Time of Indonesian Soybean towards the Production Result to Meet the Demand, International Journal of Mechanical Engineering and Technology, 9(8), 2018, pp. 1022–1028.

http://www.iaeme.com/IJMET/issues.asp?JType=IJMET&VType=9&IType=8

# **1. INTRODUCTION**

Adequate food supply, including soybeans, is the right of every citizen to be fulfilled by the government (Harsono 2008, Agricultural Research and Development Agency 2011). Domestic soybean production should be maximized to meet the needs (Suyono, 2013). Increasing soybean production to self-sufficiency level is not impossible (Supadi, 2008); the shortage of soybean production needs should be solved immediately through efforts to increase non-import production which drains the country's foreign exchange (Harsono, 2008). Superior variety of varieties can be planted throughout the year with intercropping systems tested from Sumatra to Papua (Sinar Tani 2013: Suyono 2013). During this time, domestic

1022

editor@iaeme.com

Analysis of Variety and Planting Time of Indonesian Soybean towards the Production Result to Meet the Demand

soybean farmers have been planting only once in a year, namely June or July, or at most twice a year in June and December (Nelly, 2016). One indicator to increase domestic soybean production is growing soybean throughout the year (Nelly, 2017). Therefore a research by planting outside the planting time of farmers needs to be done.

#### **2. THEORITICAL REVIEW**

The experiments done with the guidance from soybean experts obtained from the Village Integrated Service Unit (UPTD) Bangsal Sari, Jember Regency, East Java (Djoko, 2015, Imam, 2015). East Java is the largest producer of soybeans compared to other regions throughout Indonesia (Nelly, 2017). The cultivation method chosen by the researchers is the simplest method and easy to apply, considering this research uses farmers who are not soybean growers. Similarly, if the result of this study is significant as stated by researchers in previous studies that planting soybean does not need to pay attention to soil conditions and time, then the results will be applicable as additional information for other researchers and those who need. In that case the purpose to fulfil the demand of soybeans can meet the quality and reasonable price. The data analysis used experimental design theory of 2 factors with random model (Suwanda, 2015). Variety is A factor and Planting time is B factor, as the independent variable. The dependent variable is the result of production or the harvest. The reason of using a random model is because soybean cultivation in domestic production has many varieties, then the conclusion of the research can be applied to all varieties of soybean domestic production includes 5 varieties of soybeans tested. Likewise the planting time is not only for the three months tested only, but can be applied to other months; therefore it will help reducing soybean deficiency.

#### **3. RESEARCH METHODOLOGY**

The experiment was conducted with experiment that used five seeds and three planting time. Data processing and analysis used 2-factor experimental design with random model. After doing the cultivation, the next step is soil processing, planting seeds, fertilization, irrigation, harvest maintenance, threshing and cleaning then record the results.

#### 4. RESULTS ANALYSIS

The process of planting domestic soybean is based on the theory and guidance of soybean research and development unit, Ministry of Agriculture especially food crops, Jember regency, East Java province, Indonesia, BBPP 2015 (Agricultural Training Center) Ketindan, Djoko Sumianto, Agribusiness Technical Training Soy.

Also Imam Sutrisno, 2015, Soybean Cultivation Training, Research Institute of Aneka Beans and Tubers, Balit Kabi. The data processing and analysis obtained from the reference of Suwanda, Experimental Design For Scientific Research, ALFABETA, Bandung, 2017.

1023

The results of the experiment are obtained as seen in Table 1.

Planti ng time (B)	Raja Basa	Mu tiara 1	Dena 1	Dega 1	Grobogan	amount	average
March	2,2 2,9 2.4	2,7 3,7 3,5	2,2 2,5 2,4	3,0 2,9 3,4	3,2 2,9 2,8		
amount	7,5	9,9	7,1	9,3	8,9	42,7	
Average	2,5	3,3	2,4	3,1	3,0	-	2,9
April	2,4 3,4	2,7 2,8 2,5	2,1 2,5 2,2	2,8 3,4 3,2	2,8 3,0 2,0		
amount	<u> </u>	9	6,9	9.4	8.7	43	
Average	3	3	2,3	3,1	2,9	-	2,9
May	2,1 3,2 2,7	2,4 2,8 3,2	2,1 2,2 2,7	2,9 2,8 2,9	2,8 3,0 3,0		
Amount	8	8,4	7	8,6	8,8	40,8	-
Average	2,7	2,8	2,3	2,9	2,9	-	2,7
Total amount	24,5	27,3	27,3	27,3	26,4	132,8	
Average	2,6	3,0	3,0	3,0	2,9		2,9

Table 1 Results of Treatment of 5 Varieties of Indonesian Soybean and 3 Planting Time (Ton/ha)

#### **5. HYPOTHESIS**

There is no difference in the use of seed type, planting time and interaction of seed type and planting time toward production results.

: Varians:  $(\sigma^2 A) = 0$ ;  $(\sigma^2 B) = 0$ ;  $(\sigma^2 AB) = 0$ 

(There are no different variant for varieties, planting time and interaction A and B to production results)

# 6. DATA PROCESSING

$$\sum Y^2 = (2,2)^2 + (2,9)^2 + \dots + (3,0)^2 + (3,0)^2$$
  
= 372,15  
Ry = (132,8)<sup>2</sup> / 5x3x3 = 391,9  
Ay = (42,7)<sup>2</sup> + (43)<sup>2</sup> + (40,8)<sup>2</sup> / 3x3 - 391,9  
= 167,8  
By =(24,5)2 + (27,3)2 + (27,3)2 + (26,4)2 / 5 x 3 - 391,9 = - 305,42  
Jab =1/3 {(7,5)2 + (9,9)2 + ... + (8,6)2 + (8,8)2/3 - 391,9 = -32,21  
ABy = - 32,1 - 167,8 - (- 305,42) = 105,52  
Ey = 372,15 - 391,9 - 167,8 - (-305,42) - (105,52) = 12,7

editor@iaeme.com

# 7. DATA ANALYSIS

For the random model, F value (sample) / treatment is obtained by the following formula: : A = A/AB; B = B/AB; AB = AB/E

Variation source	Degree of freedom (dk )	Sum of square s (JK)	Averag e Sum of squares (RJK )	F calculation
Average	1	391,9	391,9	
OI TDEAT				
MENT.				
(Variety)				
(variety)	4	167,8	41,95	0,79
(planting				
time)	2	305.42	-152.7	-2.89
Ŧ.	-	505,12	102,7	-2,09
Inter-				
action	2	105,52	52,76	125,61
Standar				
Error	30	12,7	0,42	

Table 2 Variance Analysis of 5 Indonesian Soybean Varieties and Planting Time

With 95% confidence level, it is obtained:

F <sub>0,05</sub> Variety	(4,30) = 2,69
F <sub>0,05</sub> Planting time	(2,30) = 3,32
F <sub>0,05</sub> Interaction	(2,30) = 2,32

F calculation <F table, then the hypothesis is accepted, meaning there is no effect of production result because of the Indonesian soybean varieties, similar to the planting time. For Interaction, F calculation> F table, then the hypothesis is rejected, it means there is influence in production result because of the interaction between soybean varieties of Indonesia and Planting Time.



Figure 1 Average production of 3 planting time

From Figure 1 it can be seen that April reach the highest average production of 43 tons/ha, while March and May each produced average production of 42.7 tons/ha and 40.8 tons / ha.

1025

# 8. DISCUSSION

Table 2 showed that F calculation <F table, therefore the hypothesis is accepted. There is no effect to production result because of variety, same goes to planting location. F calculation> F table, then the hypothesis is rejected, meaning there is influence of production result because of the interaction between variety and plant location. This corresponds to the fact that each type of seed has its own profile which has been discovered by its breeder planted at the breeding site in the usual planting time of soybean farmers. For the interactions, it is proven that there is a difference in the amount of production in each planting time. The results of the research proved that soybeans can grow and can be productive if planted in other locations as well when planted in different month than usual, for instance June or July in Jember and November or December in Banyuwangi (Nelly, 2016), this is being the novelty of the researcher because previous farmers worried about their soybean for not growing or producing.

# 9. CONCLUSION

- 1. Indonesian soybean varieties do not affect the results of production obtained.
- 2. Planting time does not affect the results of production obtained.
- 3. The interaction between varieties and planting times greatly affects the production results obtained.
- 4. Planting time in May produced the highest production of 2.7 tons/ha, while March and April each produce 2.5 tons/ha
- 5. In March, the Mutiaral variety produced the highest average production of 3.3 tons/ha, in April the Dega 1 variety was 3.1 tons / ha and in May the Dega 1 and Grobogan varieties were 2, 9 tons/ha
- 6. Each variety has its own profile.

# REFERENCES

- Atman. 2009. Strategi Peningkatan Produksi Kedelai di Indonesia. Jurnal Ilmiah Tambua, 8(1), ISSN 1412-583839.
   Atman 2009 (Strategy in Increasing Soybean Production in Indonesia), Tambua Scientific Journal, 8(1), ISSN 1412-583839.
- Badan Pusat Statistik. 2014. BPS 2013.Jakarta.Statistics Central Bureau 2014 (BPS 2013),Jakarta.
- Balitkabi. 2010. Deskripsi Varietas Unggul Kacang-kacangan dan Umbi-umbian. Malang
  Balai Penelitian Kacang-kacangan dan Umbi-umbian, Balitkabi.
  Balitkabi 2010 (Description of Superior Varieties of Beans and Tubers), Research
  Institute for Pulses and Tubers, Balitkabi, Malang.
- BKPM. 2009. Budidaya kedelai di Propinsi Jawa Timur. Jawa Timur
  BKPM 2009 (Coordinating Agency for Capital Investment), Soybean Cultivation in East Java Province
- [5] Direktorat Jendral Tanaman Pangan. 2010. Road Map peningkatan produksi kedelai tahun 2010 2014. Jakarta : Kementan.

Directorate General of Food Crops 2010 (Road Map Soybean Production Increase in 2010-2014), Jakarta: Kementan

[6] Direktorat Jendral Tanaman Pangan.2014. Program dan kegiatan pembangunan pertanian tanaman pangan 2015-2019.

Directorate General of Food Crops 2014 (Program and activities of Food Crops development in 2015 – 2019).

- [7] Djoko Sumianto. 2015. Diklat Teknis Agribisnis Kedelai. Ketindan: Balai Besar Pelatihan Pertanian.
  Djoko Sumianto 2015 (Technical Training of Soybean Agribusiness), Ketindan: Center of Agricultural Training.
- [8] Harsono. 2008. Strategi pencapaian swasembada kedelai melalui perluasan areal tanam di lahan kering masam.Bogor : IPTEK Tanaman Pangan, 3(2).
  Harsono 2008 (Strategy for Achieving Soy Self-Sufficiency through Expansion of

Planting Areas on Dry Dies), Science of Food Crops, 3 (2), Bogor, 2008

- [9] Hassan, F.S.C., Fakheri,B. and Sattari, A. 2014. Review: Breeding for Resistance to Soybean Rust. International Journal of Agriculture and Crop Sciences (IJACS), 7(6), 322-328,ISSN 2227-670X ©2014 IJACS Journal.
- [10] F.S.C.Hassan, B. Fakheri and A. Sattari, 2014 (Review: Breeding for Resistance to Soybean Rust), International Journal of Agriculture and Crop Sciences (IJACS), 7(6), 322-328, ISSN 2227-670X ©2014 IJACS Journal, 2014.
- [11] Heriyanto. 2012. Upaya percepatan respon petani dalam peningkatan kontribusi varietas unggul kedelai terhadap pendapatan daerah Jawa Timur. Jurnal Cakrawala, 6(2), Juni 2012: 144-128.
  Heriyanto 2012 (Improving farmers response in increasing contribution of superior variety)

Heriyanto 2012 (Improving farmers response in increasing contribution of superior variety towards East Java regional income), Cakrawala journal, 6(2), Juni 2012:144-128.

[12] Mahasi, J. M., Mukalama, J., Mursoy, R.C., Mbehero, P. and Vanluwe, A. B. 2011. A sustainable approach to increased soybean production in western Kenya, African Crop Science Conference Proceedings, 10, d115 – 120, ISSN 1023-070X/2011.
 Mahasi, J. Mukalama, R.C. Mursoy, P. Mbehero and A. B. Vanluwe, 2011 (A sustainable approach to increased soybean production in western Kenya), African Crop Science

Conference Proceedings, 10, d115 – 120, ISSN 1023- 070X/2011.

[13] Nelly, B., Praktikto, Soedjito, S., dan Purnomo, B. S. 2015. Analisa Faktor Produksi Untuk Mencukupi Ketersediaan Kedelai Nasional. Prosiding Seminar Nasional Sains dan Teknologi. SAINTEK, Vol.1ISSN 2407-4845. Malang : Universitas Brawijaya, Teknik Mesin.

Nelly B, Praktikto, Soedjito S and Purnomo B.S. 2015 (Production Factor Analysis To Suffer National Soybean Availability, Proceedings of National Seminar on Science and Technology, SAINTEK- 'VOL.1, April 2015, ISSN 2407-4845), Journal on line, mesin.ub.ac.id/saintek, ISSN Online 2407-5329

- [14] Nelly, B., Praktikto, Soedjito, S., dan Purnomo, B. S. 2016. National Soybean Production Enhancement Strategy for Adequate Availability to Achieve Self Sufficiency, ARPN Journal Agricultural and Biological Science (JABS), Vol. 11, No. 10.
- [15] Nelly, B., Praktikto, Soedjito, S., dan Purnomo, B. S. 2017 (Determining Factor and Indicator for Alternatif National Soybean Enhancement Production Model), Journal of Engineering Science and Technology (JESTEC), Volume 12 No. 2.
- [16] Nelly B., Putu, Sonny, Strategy Pengendalian Persediaan Kedelai Dalam Negeri Guna Mencapai Swasembada, Jurnal PASTI Volume XI No. 2, 195 199 (2017).
  Nelly B., Putu, Sonny (Strategy of Maintaining Domestic Soybean Supply to reach Self-Sufficiency), PASTI journal Volume XI No. 2, 195 199 (2017).
- [17] Sinar Tani. 2013. Pengembangan Kedelai di Kawasan Hutan Sebagai Sumber Benih, Agroinivasi, 15(3740). Balitbang Pertanian.
   Sinar Tani 2013 (Development of Soybean in Forest area as Seed Source, Agroinivasi, 15(3740)), Research Insitute of Farming.

- [18] Supadi. 2008.Menggalang partisipasi petani untuk meningkatkan produksi kedelai menuju swasembada. Jurnal Litbang Pertanian Bogor.
  Supadi 2008 (Raising Farmers' Participation to Increase Soybean Production to Self-Sufficiency), Bogor Agricultural Research Journal.
- [19] Sudaryanto, T. dan Swastika, D. K. S. 2007. Kedudukan Indonesia Dalam Perdagangan Internasional Kedelai, Dalam Sumarno et al (Eds Kedelai : Teknik produksi dan Pengembangan, p. 28-44). Bogor: Puslitbang Tanaman Pangan.
   Sudaryanto, T. and D.K.S. Swastika 2007 (Indonesia's Position in International Trade of Soybean, p. 28-44 In Sumarno et al Eds Soy: Production and Development Technique), Research Center for Food Crops, Bogor
- [20] Suwanda.2017. Desain Eksperimen Untuk Penelitian Ilmiah.Bandung : ALFABETA. Suwanda 2017 (Experimental Design for Scientific Research), ALFABETA, Bandung.
- [21] Suyono.2013.Tawarkan Solusi Untuk Swasembada Kedelai.Diakses pada 20 September 2013, dari www.uhej.ac.id/berita/akademik2013.
  Suyono 2013 (Offering A Solution for Self-sufficiency. Accessed in September 20th,2013), www.uhej.ac.id/berita/akademik2013